

1 **Interplay of hydrology, community ecology and pollutant**
2 **attenuation in the hyporheic zone**

3 Ignacio Peralta–Maraver^{*}, Julia Reiss & Anne L. Robertson

4 Department of Life Sciences, Roehampton University, London, UK.

5 [*nacho.peralta@roehampton.ac.uk](mailto:nacho.peralta@roehampton.ac.uk)

6 *orcid.org/0000-0002-3467-7664

Summary

1) In this review, we describe the hierarchical interplay of hydrology, hyporheic ecology and transformation of nutrients and pollutants in the hyporheic zone (HZ). The interchange of water between the surface-subsurface generates the hyporheic hydrodynamics: the engine that drives the ecological functioning of the HZ. The magnitude and direction of hydrological fluxes in the HZ follow complex spatial patterns, strongly influenced by the temporal dynamics of surface flow in rivers.

2) The direction and magnitude of hydrological fluxes also shapes the structure of hyporheic communities (hyporheos). During surface disturbances such as flooding or drought, benthic organisms may also use the HZ as a refuge, although the importance of this role is debated.

3) Streambed organisms differ in their ability to colonise the HZ depending on the biological traits they possess. The reduction in oxygen concentration and pore size with increasing depth imposes a limit on the distribution of macroinvertebrates, which are replaced by a suite of smaller organisms (meiofauna and protists) at deeper sediment layers. Therefore, a concomitant reduction in net biomass and productivity might be expected along a vertical gradient. However, only a few studies have assessed the contribution of the hyporheos to whole system production, and they have focused only on the fraction of relatively large organisms.

4) The bioreactor ability of the HZ to transform nutrients and pollutants is an important ecosystem service sustained by the life activities of hyporheos. Biofilms have the key role in this process due to their capacity to metabolize a wide range of dissolved

compounds, including emerging pollutants. However, the residence time of water in pore sediments (resulting from hyporheic hydrodynamics) and the rest of the community (constantly reworking the sediments and grazing biofilms) are indirectly involved.

5) In order to understand how hyporheic hydrodynamics may determine the ecology of hyporheos, and its implications for the bioreactor functioning of HZ, future research in this system should combine modern and truly interdisciplinary approaches at the interface of hydrology, geomorphology and ecology.

Key words: macroinvertebrates, meiofauna, biofilms, micropollutants, bioremediation, ecosystem services.

Introduction

In most lotic systems, the surface water of the open channel is connected to groundwater systems via the riverbed sediments. As a result, there is a bi-directional exchange between the groundwater and the surface along the continuum of stream and rivers (Bencala 1993). The volume of sediments in which stream water mixes with groundwater is known as the hyporheic zone (HZ). HZ functioning in the context of the whole-river ecosystem has been studied by researchers belonging to many different disciplines and as a result selecting a single inclusive definition for the HZ is difficult (Bencala 2000). Traditionally its definition has depended on the discipline-specific interest in hyporheic processes (Tonina & Buffington, 2009; Table 1). For example, in Geochemistry, the HZ is defined as the volume of sediment containing a specified percentage of surface water, while in Biology it is described as the volume of sediments housing a characteristic hyporheic community. Even the vertical extent of the HZ is not unanimously agreed. These differences in definition and extent have important implications for study methodology. However, the fundamental concept behind all definitions is that water exchanges between the open channel and the ground water systems.

Recently, Ward (2016) proposed a more flexible and cross-disciplinary definition (Table 1). A key idea from this definition of HZ is the importance of the temporal scale relevant to the processes of interest. In fact, flow paths and the rates of water exchange through the HZ are strongly influenced by the temporal dynamics of surface flow in rivers. This is especially evident on a seasonal scale. Despite the constant dynamism to which rivers are subjected, seasonality may result in a set of drastic changes in water flow conditions

(Gasith & Resh 1999) and determine the location and extent of HZ (Wondzell 1993). Nonetheless, the HZ buffers the amplitude of this variation, acting as a potential refuge of riverbed biota during adverse conditions (See Stubbington 2012). This has important implications for variation in the composition and abundance of organisms throughout the year (Stubbington et al. 2009). The HZ harbours diverse and productive communities whose distribution and composition is strongly correlated with the direction and magnitude of hydrologic fluxes (Stanley and Boulton 1993, Olsen and Townsend 2003). These hyporheic communities or hyporheos (*hypo* = under, *rheos* = river) are composed of microbial biofilms (bacteria and fungi existing in an exocellular matrix, Singer et al. 2006), protists (mainly ciliates, flagellates and amoebae) and invertebrates. These groups differ notably in their biological traits and ability to colonize the riverbed, shaping the budget of biomass and secondary production in the HZ.

The HZ is a mechanical filter mediated by the pore space of sediments and water flows, a biochemical filter controlled by biological and chemical processes, and a photic filter (Boulton et al. 2010). As a result, the HZ provides an important ecosystem service acting as a true bioreactor (hyporheic bioreactor, Table 1) with an impressive self-purification capacity in lotic systems, and a barrier against contamination of aquifers, which are essential in the supply of water for human consumption (Lewandowski et al. 2011). Thus the HZ of streams and rivers has a critical role in the flows of biomass and energy, cycling of nutrients and pollution attenuation (McClain et al. 2003, Smith et al. 2009, Robertson & Wood 2010, Boulton et al. 2010). A large body of literature describes the nitrogen, phosphorus and organic carbon attenuation in the HZ of streams and rivers (i.e. Harvey et al. 2013, Aubeneau et al. 2015, Stegen et al. 2016, Liu et al. 2017). However, there are few data assessing the

fate and removal rates of the emerging micropollutants (Table 1) in lotic systems (Lewandowski et al. 2011, Köhler & Triebkorn, 2013), making understanding about the processing of these compounds by the bioreactor a remaining challenge in ecology.

The role of the hyporheic bioreactor in the whole river system might be seen, in an analogous way, as the ‘rivers liver’ (Fischer et al. 2005). HZ has an important role on the production, metabolisms, interchange and transformation of dissolved compounds, and health of the whole ecosystem. Here we describe the hierarchical relationship between hyporheic hydrodynamics, community ecology, and pollutant attenuation of the HZ. These subjects have been mainly assessed separately in discipline specific studies but they are intimately connected and together drive the functioning of the hyporheic bioreactor.

Hyporheic hydrodynamics: the motor of hyporheic zone ecology

The water interchange between the open river channel and a groundwater system generates the hyporheic hydrodynamics and strongly influences the whole ecosystem by determining the transport of solutes between compartments (Ward et al. 2012) and the chemistry of stream water (Duff and Triska 1990, Bencala et al. 1993). Streambed sediments are a porous medium through which exchange of water occurs. Thus, the hyporheic exchange and the flow paths throughout sediments could be theoretically studied by applying Darcy’s Law, as the product of hydraulic gradient and hydraulic conductivity values (Jones et al. 1996). However, there are a wide range of additional factors, such as the composition of sediment and the rate of mixing that act at different scales and also influence hyporheic hydrodynamics (Larkin & Sharp 1992, Hakenkamp et al. 1993, Boulton et al. 1998,

Buffington & Tonina 2009). Therefore, hyporheic hydrodynamics do not show a uniform pattern along the river. The flow between surface and groundwater follows complex dynamics (Rutheford & Hynes 1987), in which hydrologic up-welling (UW) and downwelling (DW) zones occur alternately (White 1990). Thus, as a result of this mosaic of physical features, sediment conditions may change substantially even at a centimetres scale (Boano et al. 2014). In addition, temporal dynamics from daily to seasonal processes result in re-adjustments in hyporheic flow through time (Wondzell 2011).

Streambed topography from sediment-scale (little dunes or pebbles) to larger geomorphologic features (i.e., riffle-pool sequence, steps), is the primary control on hyporheic exchange (Maddock et al. 1995, Vallet et al. 1996, Dahm et al. 1998, Calver 2001, Ward et al. 2012, Gomez-Velez et al. 2014). These features act as obstacles to the water flow along the open channel, extending the HZ both vertically and laterally from the stream (Harvey & Bencala, 1993), and generating sequences of DW and UW zones (Savant et al. 1987, Hendriks & White, 1991, Harvey & Bencala 1993), for example, within a single stream riffle, surface water enters the HZ at the beginning of the riffle (DW zone) and returns to the open channel at the end of it (Hendrik & White 1991, Hendriks 1993, Evans & Petts 1997).

Sediment grain size is another factor influencing hyporheic exchange (see Table 2 for a definition of terms). This is generally faster in headwater streams, which typically have shallow and steep stretches with cobble- and gravel-bed sediments (more porous). Hyporheic exchange is progressively slower, deeper and more complex as riverbed sediments become finer (less porous) in slower flow zones (Buffington & Tonina 2009). Nevertheless, quantifying this exchange is complicated because streambed materials range from relatively

homogeneous, to cases where the range of sediment sizes (and therefore the hydraulic conductivity) exceeds six orders of magnitude (Calver et al. 2001).

In recent years a wide range of available sampling techniques have been developed to determine the heterogeneous interactions between ground water and surface water (Kalbus et al. 2006). These methods range from direct measurements of water flux across the groundwater–surface water interface (i.e. seepage meter, Lee 1977), to indirect techniques such as heat tracer methods, mass balance approaches or mathematical modelling (Kalbus et al. 2006). Until recently these were almost exclusively employed in the field of hydrology and engineering (Boulton et al. 2010). However, understanding the importance of hyporheic hydrodynamics as a controlling factor of hyporheic communities and biochemical processes has led to increased implementation of hydrodynamic analysis in recent ecological studies of the HZ (Boulton 1993, Hendricks 1993, Standley and Boulton 1993, Schmid–Araya 1998, Kasahara et al. 2009, Miyake & Shigeru 2002, Malard et al. 2003, Davy–Bowker et al. 2006, Robertson & Wood 2010). One interesting strategy to address these issues, which has been widely used in hydrological studies, is to implement time–series analysis of streambed profiling–thermal records (i.e. Hatch et al. 2006, Keery et al. 2007, Irvine & Lautz 2015, Irvine et al. 2015). These methods are based on quantifying changes in phase and amplitude of temperature variations between pairs of subsurface sensors through depth (Hatch et al. 2006). Nevertheless, results exclusively from these methods are often limited and contradictory (Shanafield et al. 2011, Briggs et al. 2014).

Seasonality in flow exchange and its effect on riverbed communities

Stream and rivers are dynamic ecosystems par excellence. The incessant open channel flow produces continual movements of the substrata, changes to the streambed topography and reorganization of the channel boundaries morphology. In this manner, the temporal dynamics of surface flow in a lotic ecosystem is closely related to the spatial heterogeneity of flow paths and rate of exchange through the HZ at all scales. The multiple temporal and spatial scales and the rate of exchange collectively define the hyporheic residence time of water (Buffington & Tonina 2009). The residence time is an important property of the HZ, because most of the biogeochemical processes that occur in sediments depend on the rate of water flow through them (Mulholland & De Angelis 2000, Duff & Triska 2000). Furthermore, the temporal dynamic that affects the hydrological exchange also produces fluctuations of HZ boundaries (Gibert et al. 1990) and this variation of the HZ size determines its influence on both the surface open channel and the underlying groundwater (Vervier et al. 1992).

Seasonality in rivers is an extreme example of the temporal variation of the open channel discharge (e.g. due to snowmelt or the alternation between dry and rainy seasons). Accordingly, these changes alter the flow exchange patterns in the HZ (Kalbus et al. 2006) and may act as disturbances for sediment organisms (Robertson et al. 1995, Townsend et al. 1997, Robertson 2000). These potential disturbances may be reduced in the HZ due to its ability to maintain humidity after surface drying and remain stable during floods (Boulton et al. 1998). Thus the HZ might serve as a refuge for the local biota during disturbances events enabling recolonization of the surface once the disturbance ends (Williams & Hynes 1974, Dole-Olivier 2011). The HZ can also act a refuge for the early instars of some macroinvertebrates due to the more stable environmental conditions and reduced predator pressure (Williams 1984). Nevertheless, the importance of the HZ as a refuge is debated

(Robertson & Wood 2010), because some studies found no evidence of HZ refuge use by aquatic invertebrate fauna (Boulton et al. 2002, Olsen & Towsned 2003, James et al. 2008). In contrast, the importance of the HZ as a refuge might be more evident in seasonal intermittent streams. These systems are common worldwide and support diverse communities of aquatic organisms including many taxa that survive in dry riverbeds and/or rapidly recolonize when water returns (Stubbington & Datry 2013, Datry et al. 2014). Indeed, the influence of drought may be even more intense in streams that lack a marked seasonality (unpredictable intermittent streams, López-Rodríguez et al. 2012). This is the case for some Mediterranean streams with supra-seasonal drought, where many organisms that survived in the HZ during the dry season recolonized the stream during the first month of the wet season (López-Rodríguez et al. 2012). Use of the HZ as a refuge is not exclusive to large biota, it occurs across a wide range of organism size. Febria et al. (2012) observed that biofilms also use hyporheic sediments as a refuge from desiccation, mainly transported by hydrological pathways through the sediments. During periods of drought, the HZ supports bacteria associated with the infiltration of water and the creation of microhabitat in the sediment; when interstitial pore spaces become filled with water during flood events, HZ and the surface become connected allowing bacteria recolonization (Febria et al 2012).

Another factor that markedly affects the hydrology of streams and rivers is the seasonal change of in-stream vegetation cover. In-stream macrophytes are typically abundant in many lotic ecosystems during spring and summer; altering river flow and trapping sediments (Champion & Tanner 2000, Dodds & Biggs 2002). In-stream vegetation may reach from 0% to over 70% of spatial coverage between winter and summer in European rivers (Cotton et al. 2006). This increase in macrophytes is coupled with a drastic reduction

of the open channel flow velocity and the deposition of fine sediments (Cotton et al. 2006). Nevertheless, to our knowledge, there is no published research that assesses the effect of in-stream macrophyte dynamics on hyporheic flow and hyporheic communities. In addition, daily fluctuations in stream flow may also be caused by evapotranspiration of vegetation (including riparian vegetation) and it has been hypothesized that this transpiration enlarges hyporheic flow paths during the day and decreases them at night (see Wondzell et al. 2010).

Hyporheic zone as a budget of biomass and production in stream and rivers

Defining system boundaries is an important aspect of the study of ecological processes (Smock et al. 1992). Streams and rivers have been viewed traditionally as having three interactive spatial compartments: open channel and benthic zone (BZ), HZ and riparian zone (Cummins et al. 1983, Ward et al. 1989) and each compartment could play a different role depending on the ecological process under study. Secondary production is a useful measure of the energy flux (as biomass) produced by heterotrophic organisms over time and space (Benke & Huryn, 2007). However, there are only a few studies which have determined the relative contribution of the HZ compartment to whole system production (i.e. Smock et al. 1992, Collier et al. 2004, Wright–Stow et al. 2006, Reynolds & Benke 2012). Most previous studies have defined the top 5–10 cm sediment layer as the benthic zone (BZ), and lower depths as HZ despite its proposed biological definition (Table 1). In order to accurately define the limits of the HZ compartment, a small-scale approach across a depth gradient is needed.

Some studies have shown that invertebrate assemblages comprising a suite of relatively few and large individuals near the surface are replaced by numerous but small-bodied organisms with increasing depth (Schmid–Araya 1994, Stead et al. 2004). This is because the hyporheos differ in their ability to utilize the HZ depending on their biological attributes (Robertson & Wood 2010, Descoux et al. 2014), and their ability to penetrate into the HZ (Nogaro et al. 2009). The reduction in oxygen concentration and pore size due to sediment agglomeration along the depth gradient (Fig 1) limits the distribution of large macroinvertebrates with higher metabolic rates (Maridet & Philippe 1995, Strayer et al. 1997). As a result, the density of meiofauna (microscopically small metazoans) and protists should increase with depth (Fig 1). In fact, the reduction in density of large organisms through depth has been broadly reported as a general pattern in studies of riverbed communities (i.e. Dole–Olivier et al. 1994, Marchant, 1995, Maridet & Philippe 1995, Davy–Bowker et al. 2006, Pacioglu & Robertson 2017). Accordingly, it might be hypothesized that the depth gradient, as a set of different physicochemical factors, is also a key variable causing the decline of biomass and secondary production of riverbed systems. Metadata analysis of invertebrate communities from different river systems corroborates this prediction, showing a negative and significant effect of depth both on biomass and secondary production (Fig 2, Explanation of these analyses are available in appendix 1). However, despite its significance, the regression model explains only a small part of the observed variation (marginal $R^2 = 11\%$, and conditional $R^2 = 54\%$, see appendix 1) and so other variables must be important. This meta-analysis is limited to macroinvertebrate communities because studies of changes in biomass and secondary production along the depth gradient have focused exclusively on large size organisms ignoring a great fraction of meiofauna and protozoa.

Remembering that hyporheos differ in their ability to penetrate into the HZ depending on their size, we expect a significant interaction between depth and size group (flagellates, ciliates, meiofauna and macroinvertebrates). Including these groups and their interaction with depth, hydrology and sediment characteristics in future research studies would notably improve predictive modelling and compartment comparisons. This is of particular interest because it has been proposed that the bioreactor ability of HZ is sustained and maintained by diverse and active hyporheic communities (Krause et al. 2009). Accordingly, it might be predicted that hot spots of nutrient and pollutant transformation may coincide with areas containing higher biomass and secondary production rates. It could also be expected that the role of different organisms in the bioreactor capacity would vary following the depth gradient. For example, bioturbation and bioirrigation resulting from life activities of relatively large burrowers (such as Chironomidae larvae, Ephemeridae nymphs or Oligochaeta), would be more important in the benthic zone and upper layers of the HZ than in deeper levels. These processes promote sediment permeability, respiration of freshwater sediments and bacterial activity (Betrics and Ziebis 2009, Hölker et al. 2015, Baranov et al. 2016), and so have a great impact on water biogeochemistry (Morad et al. 2010).

The hyporheic bioreactor

Flow exchange and pore water chemistry of HZ can be also affected by anthropogenic activity, typically with negative effects on ecosystem health. A common alteration that occurs in rivers across the world is the artificial water input from Wastewater Treatment Plants (Carey & Migliaccio 2009) where many rivers undergo daily inputs of nutrient (i.e. organic

carbon, nitrate, phosphate) and other pollutants (i.e. pesticides) as a result of human activities (Boyer et al. 2006, Mulholland et al. 2008). However, once these compounds penetrate into the sediments as a consequence of the hydrological patterns, they may be transformed into oxidized or reduced substances by metabolic reactions, mediated by active and productive hyparctic communities (Krause et al. 2009, Bardini et al. 2012, Sánchez-Perez et al. 2013).

Accordingly, the HZ acts as a true bioreactor with water purifying ability and microbial biofilms play an important role. Hyporheic biofilms are dominated by highly diverse bacteria and archaea communities embedded in the same matrix of polysaccharides (Batin et al. 2016). This results in the coexistence of a great range of operational taxonomic units (Zeglin 2015), diverse metabolic capabilities (Singer et al. 2010, Battin et al. 2016) and sites of high enzymatic activity (Romaní et al. 2008). Thus, in stream biofilms are also important components of the global biochemical fluxes of carbon, nitrogen and phosphorous (Mulholland et al. 2008, Battin et al. 2008, Boano et al. 2014). The supply of nutrients is assumed to be a limiting factor in determining the biomass, activity, and physiology of subsurface microbial communities (Bengtsson 1989). Thus bacterial biomass and metabolic activity should be significantly greater under situations of higher input of DOC (Foulquier et al. 2011). In addition, water is pumped in and out of the HZ and riparian zones on a daily cycle because water stage variation, generates large hydraulic gradients and enhanced mixing in highly regulated rivers (Gerecht et al. 2011). These daily fluctuations of river-stage stimulate bacterial respiration and organic carbon turnover (Stegen et al. 2016).

Micropollutants, the new challenge for the hyporheic bioreactor

Recent research has shown that nitrogen, phosphorus and organic carbon are important pollutants in the HZ and aquatic ecosystems generally (i.e. Lewandowski & Nützmann, 2010, Bardini et al. 2012, Harvey et al. 2013, Maazouzi et al. 2013). However, surface water systems and their interactions with groundwater systems are increasingly under pressure from a new group of chemicals; the micropollutants (Langenhoff et al. 2013). The occurrence of micropollutants (such as pharmaceutical and personal care products; i.e. ibuprofen or antibiotics) due to WWTP inputs has greatly increased in stream and rivers. The concern about their presence is mainly related to potential adverse effects on environmental systems (i.e. bioaccumulation) and to human toxicology (i.e. aquifer contamination) (Hernández Leal et al. 2010, Langenhoff et al. 2013). Furthermore, the chronic low-level antibiotic exposures detected in aquatic systems acts as a selective process on bacteria communities (Hirsch et al. 1999, Yang & Carlson 2003). Thus, differential antibiotic tolerance of a bacterial community may produce a shift in the biofilms structure (composition, richness, density), affect the spatial distribution of members of the community (Roose-Amsaleg & Laverman 2016) and change the ability of biofilms to conduct ecosystem services (e.g. reducing denitrification processes due to deleterious impacts on denitrifying bacteria, Costanzo et al. 2005). However, in some cases, micropollutants can be efficiently attenuated along flow paths into the HZ (Lewandowski et al. 2011). Indeed, some of these compounds (i.e. diclofenac, bezafibrate, ibuprofen, and naproxen) are more efficiently transformed in river sediments than in wastewater treatment plants (WWTP) by biofilms (i.e. Schulz et al. 2008, Radke et al. 2009). This is mainly related to the higher diversity of microbial communities in environmental systems. In addition, water residence times in the

HZ are longer than in the open channel and surface sediments (and WWTP), allowing more efficient biodegradation processes (Lewandowski et al. 2011).

Notwithstanding the role of the biofilm is recognized in pollutant attenuation, more complex questions behind this ecological process remained unanswered (i.e. the role of the rest of the community). Furthermore, it is also important to consider the hierarchical interaction between hydrological patterns and biochemical processes in the study of the nutrient and pollutant breakdown into the HZ. Hydrodynamics throughout the sediments may induce two opposite effects on solute reactions. Higher inward water fluxes lead to a larger input of substances into the HZ enhancing reaction rates, but also the hyporheic microbiota has less time to perform biochemical reactions due to the lower residence time of the compounds in the sediments (Fig 3a) (Bardini et al. 2012). These mechanisms will become even more complex when we recall that hydraulic conductivity may also be affected by the action of the hyporheos. Growth of biofilm matrices in the sediment pores reduces permeability and increases residence times of water into the HZ (Findlay & Sobczak, 2000, Battin et al. 2003). In addition, biofilm theory holds that uptake of solutes is diffusion limited by the thickness of the biofilm polysaccharide matrix (Gantzer 1988). Before assimilation, solutes must pass first from the pore water to the biofilm surface (external mass transfer) and then through the biofilm matrix to the cells (internal mass transfer) (Battin et al. 2003). However, sediments are constantly being reworked, increasing the sediments permeability locally (Boulton, 2000). Life activities of macroinvertebrates, meiofauna and protists (e.g. ciliates and flagellates) in the HZ (digging, removing the sediments and grazing on biofilms) result in preferential flow paths, increasing biofilm surface and boosting bacterial densities (Danielopol 1976, Boulton 2000, Mermillod–Blondin et al. 2003, Battin et al. 2003), acting

as ecosystem engineers (Fig 3c, d). Thus, the net effect on breakdown rates depends on the balance between all these opposing factors (Arnon et al. 2007, Cardenas et al. 2008, Bardini et al. 2012). In this manner, mechanistic understanding of biofilms function can be acquired only through carefully designed experiments under well-defined conditions and appropriate cultivation techniques (Singer et al. 2006). Accordingly, controlled experiments are needed to explore the underlying causal mechanisms that generate the patterns seen in reach-scale descriptive surveys (Boulton et al. 1998, Olsen and Townsend 2003).

Concluding remarks

Ecological research in the HZ is still a challenge due to the great number of variables involved in its operation. Furthermore, there is a notable hierarchy of these variables, many of them being nested or correlated spatially and temporally. Its study is also becoming one of the most developed areas in freshwater science in recent years, principally due to the development and increasing accessibility of better technological approaches. Nevertheless, studies on the role of the HZ in the functioning and ecosystem service delivery of the whole river system are still maturing and remain a major research focus and challenge for future freshwater researchers (Robertson and Wood 2010). Complex and interesting questions about how pollutants and nutrients are transformed by the hyporheos during their travel throughout the HZ, or how hyporheic hydrodynamics may determine energy fluxes of these communities are still unanswered. Furthermore, future research should also consider and focus on a more thorough understanding of impacts caused by rivers regulation on hyporheic hydrodynamics (Gerecht et al. 2011). Accordingly, holistic and truly interdisciplinary approaches at the

interface of hydrology, geomorphology and ecology will be the only valid strategy to assess all these issues (Pacioglu 2010, Boulton et al. 2010, Robertson and Wood 2010).

Acknowledgements

This project and Ignacio Peralta–Maraver PhD–studies were funded by the European Union’s Horizon2020 research and innovation programme under Marie–Skłodowska–Curie grant agreement No. 641939.

Appendix

Biomass and secondary production data along depth gradient were extracted from reported values in Smock et al. (1992) and Reynolds and Benke (2012). A similar sampling method (sediment corers) and spatial scale resolution were used in both studies. Reynolds and Benke (2012) reported biomass and secondary production of the whole community of macroinvertebrates in a sandy–sediment river (Buzzards Branch). While, Reynolds and Benke (2012) measured genus–specific biomass and secondary production of chironomid larvae (Diptera) assemblage (composed by 26–31 genera) along a hyporheic gradient, comparing gravel–cobble of a high–alkalinity stream (Hendrik Mill Brach) with sandy and mud–silt habitats of a low–alkalinity stream (Payne Creek). Those chironomid assemblages were reasonably similar between studied streams. Furthermore, chironomid larvae are usually the dominant and richest invertebrate group in freshwater benthic habitats (Ferrington et al. 2008). Hence, it could be assumed as a good model group to infer general production patterns of invertebrates.

Two linear mixed effect models (LMMs) were applied to test the effect of, depth (continuous covariant) on biomass and production (responses). Responses were Log_{10} transformed to solve heterogeneity in the residuals. An Applicable Information Criterion (AIC) was then used to find the most parsimonious model by combining the fixed term (depth) and potential random effects (i.e. studied site, differences between taxa). As result, biomass and production models included depth (single covariate), study site (studied river as random intercept) and the interaction between depth and taxa (random slope) as effective parameters:

$$\text{Log}_{10}(\hat{y}_{ij}) = \beta_0 + a_{site} + \beta_1 \times \text{Depth}_j + b_i \times \text{Depth}_j + \varepsilon_{ij}$$

$$\varepsilon_{ij} \sim \text{Norm}(0, \sigma^2)$$

$$a_{site} \sim \text{Norm}(0, \sigma_{site}^2)$$

$$b_i \sim \text{Norm}(0, \sigma_i^2)$$

where \hat{y}_{ij} is the biomass or production for each taxa i at depth j (0,10...,50). Intercept of the model is given by $\beta_0 + a_{site}$ with changes randomly by a_{site} , and $b_i \times \text{Depth}_j$ represent the random variation of slope β_1 .

Model validation was applied following Zuur et al. (2009). Previous models were fitted using the restricted maximum likelihood estimation (REML) with functions *lmer* of the R package lme4 (Boulton et al. 2016, R Core Team 2016). Finally, 5000 values from the posterior joint distribution of the model parameters were simulated with the function *sim* of the R package arm (Gelman and Hill 2007). This function uses an analytical direct-simulation method with uninformative priors (Korner–Nievergelt 2015). Obtained means of the simulated values from the joint posterior distribution of model parameters were used as

estimates, and the 2.5% and 97.5% quintiles as lower and upper limits of 95% credible intervals. Finally, the marginal and conditional R^2 (as a technique to describe the predictive capacity of mixed effect models; Nakagawa & Schielzeth 2013) was calculated to assess model fit.

References

- Arnon, S., Gray, K.A., Packman, A.I., 2007. Biophysicochemical process coupling controls nitrogen use by benthic biofilms. *Limnol. Oceanogr.* 52, 1665–1671.
- Aubeneau, A.F., Drummond, J.D., Schumer, R., Bolster, D., Tank, J. L. Packman, A. I., 2015. Effects of benthic and hyporheic reactive transport on breakthrough curves. *Freshw. Sci.* 34, 301–315.
- Baranov, V., Lewandowski, J., Romeijn, P., Singer, G., Krause, S., 2016. Effects of bioirrigation of non-biting midges (Diptera: Chironomidae) on lake sediment respiration. *Sci. Rep.* 6: 27329.
- Bardini, L., Boano, F., Cardenas, M.B., Revelli, R., Ridolfi, L., 2012. Nutrient cycling in bedform induced hyporheic zones. *Geochim. Cosmochim. Acta.* 84, 47–61.
- Bates, D., Maechler, M, Bolker, B., Walker, S., 2016. lme4: Linear Mixed-Effects Models using 'Eigen' and S4. R package version 1.1–12. <https://cran.r-project.org/web/packages/lme4/index.html>
- Battin, T. J., Besemer, K., Bengtsson, M. M., Romani, A. M., & Packmann, A. I. (2016). The ecology and biogeochemistry of stream biofilms. *Nature Rev. Microbiol.* 14, 251–263.
- Battin, T.J., Kaplan, L.A., Findlay, S., Hopkinson, C.S., Marti, E., Packman, A.I., Newbold, J.D., Sabater, F., 2008. Biophysical controls on organic carbon fluxes in fluvial networks. *Nat. Geosci.* 1, 95–100.

448 Battin, T.J., Kaplan, L.A., Newbold, J.D., & Hansen, C.M., 2003. Contributions of microbial
449 biofilms to ecosystem processes in stream mesocosms. *Nature*. 426, 439–442.

450 Bencala, K.E., 1993. A perspective on stream–catchment connections. *J. North Am.*
451 *Benthological Soc.* 12, 44–47.

452 Bencala, K.E., 2000. Hyporheic zone hydrological processes. *Hydrol. Process.* 14, 2797–
453 2798.

454 Bengtsson, G., 1989. Growth and metabolic flexibility in groundwater bacteria. *Microb. Ecol.*
455 18, 235–248.

456 Benke, A.C., Huryn, A.D., 2007. Secondary production of macroinvertebrates, in: Hauer,
457 F.R., Lambet, G.A. (Eds.), *Methods in stream ecology*. Academic Press, San
458 Diego, pp. 691–710.

459 Bertics, V.J., Ziebis, W., 2009. Biodiversity of benthic microbial communities in bioturbated
460 coastal sediments is controlled by geochemical microniches. *ISME J.* 3, 1269–
461 1285.

462 Boano, F., Harvey, J.W., Marion, A., Packman, A. I., Revelli, R., Ridolfi, L., Wörman, A.,
463 2014. Hyporheic flow and transport processes: Mechanisms, models, and
464 biogeochemical implications. *Rev. Geophys.* 52, 603–679.

465 Boulton, A.J., Findlay, S., Marmonier, P., Stanley, E.H., Valett, H.M., 1998. The functional
466 significance of the hyporheic zone in streams and rivers. *Annu. Rev. Ecol. Syst.*
467 29, 59–81.

468 Boulton, A.J., 1993. Stream ecology and surface–hyporheic hydrologic exchange:
469 implications, techniques and limitations. *Mar. Freshwater. Res.* 44, 553–564.

470 Boulton, A.J., 2000. The surface macrofauna, in: Jones, J.B., Mulholland, P. J. (Eds.), *Stream*
471 *and ground water*. Academic Press, San Diego, pp. 337–361.

472 Boulton, A.J., Datry, T., Kasahara, T., Mutz, M., Stanford, J.A., 2010. Ecology and
473 management of the hyporheic zone: stream–groundwater interactions of running
474 waters and their floodplains. *J. North Am. Benthological Soc.* 29, 26–40.

475 Boulton, A.J., Findlay, S., Marmonier, P., Stanley, E.H., Valett, H.M., 1998. The functional
476 significance of the hyporheic zone in streams and rivers. *Annu. Rev. Ecol. Evol.*
477 *Syst.* 29, 59–81.

478 Boulton, A., Harvey, M., Proctor, H., 2002. Of spates and species: responses by interstitial
479 water mites to simulated spates in a subtropical Australian river. *Exp. Appl.*
480 *Acarol.* 34, 149–169.

481 Briggs, M.A., Lautz, L.K., Buckley, S.F., & Lane, J.W., 2014. Practical limitations on the use
482 of diurnal temperature signals to quantify groundwater upwelling. *J. Hydrol.* 519,
483 1739–1751.

484 Buffington, J. M., & Tonina, D., 2009. Hyporheic exchange in mountain rivers II: Effects of
485 channel morphology on mechanics, scales, and rates of exchange. *Geogr.*
486 *Compass.* 3, 1038–1062.

487 Calver, A., 2001. Riverbed permeabilities: Information from pooled data. *Groundwater* 39,
488 546–553.

489 Cardenas, M.B., 2008. The effect of river bend morphology on flow and timescales of surface
 490 water–groundwater exchange across pointbars. *J. Hydrol.* 362, 134–141.

491 Carey, R.O., Migliaccio, K.W., 2009. Contribution of wastewater treatment plant effluents to
 492 nutrient dynamics in aquatic systems: a review. *Environ. Manage.* 44, 205–217.

493 Champion, P.D., Tanner, C.C., 2000. Seasonality of macrophytes and interaction with flow in
 494 a New Zealand lowland stream. *Hydrobiologia* 441, 1–12.

495 Collier, K.J., Wright–Stow, A.E., Smith, B.J., 2004. Trophic basis of production for a mayfly
 496 in a North Island, New Zealand, forest stream: contributions of benthic versus
 497 hyporheic habitats and implications for restoration. *N. Z. J. Mar. Freshwater Res.*
 498 38, 301–314.

499 Costanzo, S.D., Murby, J., Bates, J., 2005. Ecosystem response to antibiotics entering the
 500 aquatic environment. *Mar. Pollut. Bull.* 51, 218–223.

501 Cotton, J.A., Wharton, G., Bass, J.A.B., Heppell, C.M., Wotton, R.S., 2006. The effects of
 502 seasonal changes to in–stream vegetation cover on patterns of flow and
 503 accumulation of sediment. *Geomorphology* 77, 320–334.

504 Cummins, K.W., Sedell, J.R., Swanson, F.J., Minshall, G.W., Fisher, S.G., Cushing, C.E.,
 505 Petersen, R.C., Vannote, R.L., 1983. Organic matter budgets for stream
 506 ecosystems: problems in their evaluation, in: Barnes, J.R., Minshall, G.W. (Eds.),
 507 Stream ecology. Springer US, New York, pp. 299–353.

508 Dahm, C.N., Grimm, N.B., Marmonier, P., Valett, H.M., Vervier, P., 1998. Nutrient
509 dynamics at the interface between surface waters and groundwaters. *Freshw.*
510 *Biol.* 40, 427–451.

511 Danielopol, D.L., 1976. The distribution of the fauna in the interstitial habitats of riverine
512 sediments of the Danube and the Piesting (Austria). *Int. J. Speleol.* 8, 23–51.

513 Datry, T., Larned, S.T., Tockner, K., 2014. Intermittent rivers: a challenge for freshwater
514 ecology. *BioScience*, bit027.

515 Davy–Bowker, J., Sweeting, W., Wright, N., Clarke, R.T., Arnott, S., 2006. The distribution
516 of benthic and hyporheic macroinvertebrates from the heads and tails of riffles.
517 *Hydrobiologia* 563, 109–123.

518 Descloux, S., Datry, T., Usseglio-Polatera, P., 2014. Trait-based structure of invertebrates
519 along a gradient of sediment colmation: Benthos versus hyporheos responses. *Sci.*
520 *Total Environ.* 466, 265–276.

521 Dodds, W.K., Biggs, B.J., 2002. Water velocity attenuation by stream periphyton and
522 macrophytes in relation to growth form and architecture. *J. North Am.*
523 *Benthological Soc.* 21, 2–15.

524 Dole–Olivier, M.J., 2011. The hyporheic refuge hypothesis reconsidered: a review of
525 hydrological aspects. *Mar. Freshwater. Res.* 62, 1281–1302.

526 Dole–Olivier M.J., Marmonier P., Creuzé des Châtelliers M., Martin D., 1994. Interstitial
527 fauna associated with the alluvial deposits of the Rhone River (France), in:

528 Danielopol D., Gibert J., Stanford J. (Eds), Groundwater ecology. Academic
529 Press, New York, pp. 140–176.

530 Duff, J.H., Triska, F.J., 2000. Nitrogen biogeochemistry and surface–subsurface exchange in
531 streams, in: Jones, J.B., Mulholland, P.J. (Eds.), Streams and groundwaters.
532 Academic Press, San Diego, pp. 197–220.

533 Evans, E.C., Petts, G.E., 1997. Hyporheic temperature patterns within riffles. Hydrol. Sci. J.
534 42, 199–213.

535 Febria, C.M., Beddoes, P., Fulthorpe, R.R., & Williams, D.D., 2012. Bacterial community
536 dynamics in the hyporheic zone of an intermittent stream. ISME J. 6, 1078–1088.

537 Findlay, S., Sobczak, W.V., 2000. Microbial communities in Hyporheic sediments, in: Jones,
538 J.B., Mulholland, P.J. (Eds.), Streams and Ground waters. Academic press, San
539 Diego, pp. 287–306.

540 Ferrington, L.C., Berg, M.B. Coffman, W.P., 2008. Chironomidae, in: Merritt, R.W.
541 Cummins, K.W., Berg, M.B. (Eds.), An introduction to the aquatic insects of
542 North America. Kendall/Hunt Publishing Company, Dubuque, pp. 847–975.

543 Fischer, H., Kloep, F., Wilzcek, S., Pusch, M. T., 2005. A river's liver–microbial processes
544 within the hyporheic zone of a large lowland river. Biogeochemistry 76, 349–371.

545 Foulquier, A., Mermillod–Blondin, F., Malard, F., Gibert, J., 2011. Response of sediment
546 biofilm to increased dissolved organic carbon supply in groundwater artificially
547 recharged with stormwater. J. Soils Sediments 11, 382–393.

548 Franken, R.J., Storey, R.G., Williams, D.D., 2001. Biological, chemical and physical
549 characteristics of downwelling and upwelling zones in the hyporheic zone of a
550 north-temperate stream. *Hydrobiologia* 444, 183–195.

551 Gantzer, C.J., Rittmann, B.E., Herricks, E.E., 1988. Mass transport to streambed biofilms.
552 *Water Res.* 22, 709–722.

553 Gasith, A., Resh, V.H., 1999. Streams in Mediterranean climate regions: abiotic influences
554 and biotic responses to predictable seasonal events. *Annu. Rev. Ecol. Evol. Syst.*
555 30, 51–81.

556 Gelman, A., Su, Y.S., Yajima, M., Hill, J., Pittau, M. G., Kerman, J., Zheng T., Dorie, V.,
557 2009. arm: data analysis using regression and multi-level/hierarchical models. R
558 package, version 9.01. <http://CRAN.R-project.org/package=arm>.

559 Gerecht, K.E., Cardenas, M.B., Guswa, A.J., Sawyer, A.H., Nowinski, J.D., Swanson, T.E.,
560 2011. Dynamics of hyporheic flow and heat transport across a bed-to-bank
561 continuum in a large regulated river. *Water Resour. Res.* 47, W03524.

562 Gibert, J., Dole-Olivier, M.J., Marmonier, R., Vervier, P., 1990. Surface water-groundwater
563 ecotone, in Naiman, R. J., Décamps, H. (Eds.), *The Ecology and Management of*
564 *Aquatic-Terrestrial Ecotones*. UNESCO, Paris, pp. 199–225.

565 Gomez-Velez, J.D., Krause, S., Wilson, J.L., 2014. Effect of low-permeability layers on
566 spatial patterns of hyporheic exchange and groundwater upwelling. *Water*
567 *Resour. Res.* 50, 5196–5215.

568 Harvey, J.W., Bencala, K.E., 1993. The effect of streambed topography on surface-
 569 subsurface water exchange in mountain catchments. *Water Resour. Res.* 29, 89–
 570 98.

571 Hakenkamp, C.C., Valett, H.M., Boulton, A.J., 1993. Perspectives on the hyporheic zone:
 572 integrating hydrology and biology. Concluding remarks. *J. North Am.*
 573 *Benthological Soc.* 12, 94–99.

574 Harvey, J.W., Böhlke, J.K., Voytek, M.A., Scott, D., Tobias, C.R., 2013. Hyporheic zone
 575 denitrification: Controls on effective reaction depth and contribution to whole-
 576 stream mass balance. *Water Resour. Res.* 49, 6298–6316.

577 Hatch, C.E., Fisher, A.T., Revenaugh, J.S., Constantz, J., Ruehl, C., 2006. Quantifying
 578 surface water–groundwater interactions using time series analysis of streambed
 579 thermal records: Method development. *Water Resour. Res.* 42, W10410.

580 Hendricks, S.P., 1993. Microbial ecology of the hyporheic zone: a perspective integrating
 581 hydrology and biology. *J. North Am. Benthological Soc.* 12, 70–78.

582 Hernández Leal, L., Vieno, N., Temmink, H., Zeeman, G., Buisman, C.J., 2010. Occurrence
 583 of xenobiotics in gray water and removal in three biological treatment systems.
 584 *Environ. Sci. Technol.* 44, 6835–6842.

585 Hirsch, R., Ternes, T., Haberer, K., Kratz, K.L. 1999. Occurrence of antibiotics in the aquatic
 586 environment. *Sci. Total Environ.* 225, 109–118.

587 Hölker, F., Vanni, M.J., Kuiper, J.J., Meile, C., Grossart, H.P., Stief, P., Adrian, R., Lorke,
 588 A., Dellwig, O., Brand, A., Hupfer, M., Mooji, M.W., Nitzmann, G.,
 589 Lewandowski, J., 2015. Tube-dwelling invertebrates: tiny ecosystem engineers
 590 have large effects in lake ecosystems. *Ecol. Monogr.* 85, 333–351.

591 Irvine, D.J., Lautz, L.K., 2015. High resolution mapping of hyporheic fluxes using streambed
 592 temperatures: Recommendations and limitations. *J. Hydrol.* 524, 137–146.

593 Irvine, D.J., Lautz, L.K., Briggs, M.A., Gordon, R.P., McKenzie, J.M., 2015. Experimental
 594 evaluation of the applicability of phase, amplitude, and combined methods to
 595 determine water flux and thermal diffusivity from temperature time series using
 596 VFLUX 2. *J. Hydrol.* 531, 728–737.

597 James, A.B., Dewson, Z.S., Death, R.G., 2008. Do stream macroinvertebrates use instream
 598 refugia in response to severe short-term flow reduction in New Zealand streams?.
 599 *Freshwater Biol.* 53, 1316–1334.

600 Kalbus, E., Reinstorf, F., Schirmer, M. 2006. Measuring methods for groundwater–surface
 601 water interactions: a review. *Hydrol. Earth. Syst. Sc.* 10, 873–887.

602 Kasahara, T., Datry, T., Mutz, M., Boulton, A. J., 2009. Treating causes not symptoms:
 603 restoration of surface–groundwater interactions in rivers. *Mar. Freshwater. Res.*
 604 60, 976–981.

605 Keery, J., Binley, A., Crook, N., Smith, J.W., 2007. Temporal and spatial variability of
 606 groundwater–surface water fluxes: development and application of an analytical
 607 method using temperature time series. *J. Hydrol.* 336, 1–16.

608 Köhler, H. R., Triebskorn, R., 2013. Wildlife ecotoxicology of pesticides: can we track
609 effects to the population level and beyond?. *Science* 341, 759–765.

610 Korner–Nievergelt, F., Roth, T., von Felten, S., Guélat, J., Almasi, B., Korner–Nievergelt, P.,
611 2015. Bayesian data analysis in ecology using linear models with R, BUGS, and
612 Stan. Academic Press, London.

613 Krause, S., Hannah, D.M., Fleckenstein, J.H., 2009. Hyporheic hydrology: interactions at the
614 groundwater–surface water interface. *Hydrol. Process.* 23, 2103–2107.

615 Langenhoff, A., Inderfurth, N., Veuskens, T., Schraa, G., Blokland, M., Kujawa–Roeleveld,
616 K., Rijnaarts, H., 2013. Microbial removal of the pharmaceutical compounds
617 ibuprofen and diclofenac from wastewater. *Biomed. Res. Int.* 2013, 1–9

618 Larkin, R.G., Sharp, J.M., 1992. On the relationship between river–basin geomorphology,
619 aquifer hydraulics, and ground–water flow direction in alluvial aquifers. *Geol.*
620 *Soc. Am. Bull.* 104, 1608–1620.

621 Lee, D. R., 1977. A device for measuring seepage flux in lakes and estuaries. *Limnol.*
622 *Oceanogr.* 22, 140–147.

623 Lewandowski, J., Nützmann, G., 2010. Nutrient retention and release in a floodplain's aquifer
624 and in the hyporheic zone of a lowland river. *Ecol. Eng.* 36, 1156–1166.

625 Lewandowski, J., Putschew, A., Schwesig, D., Neumann, C., Radke, M., 2011. Fate of
626 organic micropollutants in the hyporheic zone of a eutrophic lowland stream:
627 Results of a preliminary field study. *Sci. Total Environ.* 409, 1824–1835.

628 Liu, Y., Liu, C., Nelson, W. C., Shi, L., Xu, F., Liu, Y., Xu, F., Liu, Y., Yan, A., Zhong, L.,
 629 Thompson, C., Fredrikson, J.K., Zachara, J. M., 2017. Effect of Water Chemistry
 630 and Hydrodynamics on Nitrogen Transformation Activity and Microbial
 631 Community Functional Potential in Hyporheic Zone Sediment Columns. *Environ.*
 632 *Sci. Technol.* 51, 4877–4886.

633 López-Rodríguez, M.J., Peralta-Maraver, I., Gaetani, B., Sainz-Cantero, C.E., Fochetti, R.,
 634 Tierno de Figueroa, J., 2012. Diversity patterns and food web structure in a
 635 Mediterranean intermittent stream. *Int. Rev. Hydrobiol.* 97, 485–496.

636 Luo, Y., Guo, W., Ngo, H. Hao., Nghiem, L. Duc., Hai, F. Ibney., Zhang, J., Liang, S., 2014.
 637 A review on the occurrence of micropollutants in the aquatic environment and
 638 their fate and removal during wastewater treatment. *Sci. Total Environ.* 473, 619–
 639 641.

640 Maazouzi, C., Claret, C., Dole-Olivier, M. J., Marmonier, P., 2013. Nutrient dynamics in
 641 riverbed sediments: effects of hydrological disturbances using experimental flow
 642 manipulations. *J. Soils Sediments* 13, 207–219.

643 Maddok, I.P., Petts, G.E., Evans, E.C., Greenwood, M.T., 1995. Assessing river-aquifer
 644 interactions within the hyporheic zone, in: Brown, A. G. (Ed.), *Groundwater and*
 645 *Geomorphology*. Wiley, Chichester, pp. 53–74.

646 Malard, F., Ferreira, D., Dolédec, S., Ward, J.V., 2003. Influence of groundwater upwelling
 647 on the distribution of the hyporheos in a headwater river flood plain. *Arch.*
 648 *Hydrobiol.* 157, 89–116.

649 Marchant, R., 1995. Seasonal variation in the vertical distribution of hyporheic invertebrates
650 in an Australian upland river. *Arch. Hydrobiol.* 134, 441–457.

651 Maridet, L., Philippe, M., 1995. Influence of substrate characteristics on the vertical
652 distribution of stream macroinvertebrates in the hyporheic zone. *Folia Fac. Sci.*
653 *Natur. Univ. Masaryk. Brun. Math.* 91, 101–105.

654 Mermillod–Blondin, F., Gaudet, J. P., Gérino, M., Desrosiers, G., Creuzé des Châtelliers, M.,
655 2003. Influence of macroinvertebrates on physico–chemical and microbial
656 processes in hyporheic sediments. *Hydrol. Process.* 17, 779–794.

657 McClain, M.E., Boyer, E.W., Dent, C.L., Gergel, S.E., Grimm, N.B., Groffman, P.M., Hart,
658 S. C., Harvey, J.W., Johnston, C.A., Mayorga, E., McDowell, W.H., Pinay, G.,
659 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial
660 and aquatic ecosystems. *Ecosystems* 6, 301–312.

661 Miyake, Y., Nakano, S., 2002. Effects of substratum stability on diversity of stream
662 invertebrates during baseflow at two spatial scales. *Freshwater Biol.* 47, 219–230.

663 Morad, M.R., Khalili, A., Roskosch, A., Lewandowski, J., 2010. Quantification of pumping
664 rate of *Chironomus plumosus* larvae in natural burrows. *Aquatic Ecol.* 44, 143–
665 153.

666 Mulholland, P.J., DeAngelis, D.L., 2000. Surface–subsurface exchange and nutrient
667 spiralling, in: Jones, J. B., Mulholland, P. J. (Eds.), *Streams and Ground waters*.
668 Academic Press. San Diego, pp. 149–168.

669 Mulholland, P.J., Helton, A.M., Poole, G.C., Hall, R.O., Hamilton, S.K., Peterson, B.J., Tank,
 670 J.T., Ashkenas, L.R., Cooper, L.W., Dham, C.N., Dodds, W.K., Findlay, S.E.G.,
 671 Gregory, S.V., Grimm, N.B., Johnson, S.L., McDowell, L.W., Meyer, J.L.,
 672 Valett, H.M., Webster, J.R., Arango, C.P., Beaulieu, J.J., Bernot, M.J., Burgin,
 673 A.J., Crenshaw, C.L., Johnson, L.T., Niederlehner, B.R., O'Brien, J.M., Potter,
 674 J.D., Sheibley, R.W., Sobota, D.J., Thomas, S.M., 2008. Stream denitrification
 675 across biomes and its response to anthropogenic nitrate loading. *Nature* 452, 202–
 676 205.

677 Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R² from
 678 generalized linear mixed-effects models. *Methods Ecol Evol* 4, 133-142.

679 Nogaro, G., Mermillod-Blondin, F., Valett, M.H., François-Carcaillet, F., Gaudet, J.P.,
 680 Lafont, M., Gibert, J., 2009. Ecosystem engineering at the sediment–water
 681 interface: bioturbation and consumer–substrate interaction. *Oecologia* 161, 125–
 682 138.

683 Olsen, D.A., Townsend, C.R., 2003. Flood effects on invertebrates, sediments and particulate
 684 organic matter in the hyporheic zone of a gravel–bed stream. *Freshwater Biol.* 50,
 685 839–853.

686 Orghidan, T., 1955. Un nou domeniu de viata acvatica subterana: 'biotopul hiporeic'. *Bull.*
 687 *Sti. sect. Biologie si sti. Agronau. si sect. Geologie si Geogr.* 7, 657–676.

688 Pacioglu, O., 2010. Ecology of the hyporheic zone: a review. *Cave Karst Sci.* 36, 69–76.

689 Pacioglu, O., Robertson, A., 2017. The invertebrate community of the chalk stream hyporheic
690 zone: spatio-temporal distribution patterns. *Knowl Manag Aquat Ecosyst.* 418,
691 10.

692 Radke, M., Lauwigi, C., Heinkele, G., Mürdter, T.E., Letzel, M., 2009. Fate of the antibiotic
693 sulfamethoxazole and its two major human metabolites in a water sediment test.
694 *Environ. Sci. Technol.* 43, 3135–3141.

695 Reynolds Jr, S.K., Benke, A.C., 2012. Chironomid production along a hyporheic gradient in
696 contrasting stream types. *Freshw. Sci.* 3, 167–181

697 Robertson, A. L., 2000. Lotic meiofaunal community dynamics: colonisation, resilience and
698 persistence in a spatially and temporally heterogeneous environment. *Freshwater*
699 *Biol.* 44, 135–147.

700 Robertson, A.L., Wood, P.J., 2010. Ecology of the hyporheic zone: origins, current
701 knowledge and future directions. *Fundam. Appl. Limnol./Arch. Hydrobiol.* 176,
702 279–289.

703 Romani, A.M., Fund, K., Artigas, J., Schwartz, T., Sabater, S., Obst, U., 2008. Relevance of
704 polymeric matrix enzymes during biofilm formation. *Microb. Ecol.* 56, 427–436.

705 Roose-Amsaleg, C., & Laverman, A. M. (2016). Do antibiotics have environmental side-
706 effects? Impact of synthetic antibiotics on biogeochemical processes. *Environ.*
707 *Sci. Pollut. Res.* 23, 4000-4012.

708 Rutherford, J.E., Hynes, H.B.N., 1987. Dissolved organic carbon in streams and groundwater.
709 *Hydrobiologia* 154, 33–48.

710 Sánchez-Pérez, J.M., Montuelle, B., Mouchet, F., Gauthier, L., Julien, F., Sauvage, S.,
711 Teissier, S., Dedieu, K., Destrieux, D., Vervier, P., Gerino, M. Role of the
712 hyporheic heterotrophic biofilm on transformation and toxicity of pesticides.
713 *Ann. Limnol. - Int. J. Lim.* 49, 87–95.

714 Savant, S.A., Reible, D.D., Thibodeaux, L.J., 1987. Convective transport within stable river
715 sediments. *Water Resour. Res.* 23, 1763–1768.

716 Schmid–Araya, J.M., 1994. Temporal and spatial distribution of benthic microfauna in
717 sediments of a gravel streambed. *Limnol. Oceanogr.* 39, 1813–1821

718 Schmid–Araya, J. M., 1998. Rotifers in interstitial sediments. *Hydrobiologia* 387, 231–240.

719 Schulz, M., Löffler, D., Wagner, M., Ternes, T. A., 2008. Transformation of the X-ray
720 contrast medium iopromide in soil and biological wastewater treatment. *Environ.*
721 *Sci. Technol* 42, 7207–7217.

722 Shanafield, M., Hatch, C., Pohll, G., 2011. Uncertainty in thermal time series analysis
723 estimates of streambed water flux. *Water Resour. Res.* 47, W03504.

724 Singer, G., Besemer, K., Hödl, I., Chlup, A., Hochedlinger, G., Stadler, P., & Battin, T.J.,
725 2006. Microcosm design and evaluation to study stream microbial
726 biofilms. *Limnol. Oceanogr. Methods* 4, 436-447.

727 Singer, G., Besemer, K., Schmitt–Kopplin, P., Hödl, I., & Battin, T.J., 2010. Physical
728 heterogeneity increases biofilm resource use and its molecular diversity in stream
729 mesocosms. *PLoS One* 5, e9988.

730 Smith, J.W.N., Surridge, B.W., Haxton, T.H., Lerner, D.N., 2009. Pollutant attenuation at the
731 groundwater–surface water interface: A classification scheme and statistical
732 analysis using national–scale nitrate data. *J. Hydrol.* 369, 392–402.

733 Smock, L.A., Gladden, J.E., Riekenberg, J.L., Smith, L.C., Black, C.R., 1992. Lotic
734 macroinvertebrate production in three dimensions: channel surface, hyporheic,
735 and floodplain environments. *Ecology* 73, 876–886.

736 Stanley, E.H., Boulton, A.J., 1993. Hydrology and the distribution of hyporheos: perspectives
737 from a mesic river and a desert stream. *J. North Am. Benthological Soc.* 12, 79–
738 83.

739 Stead, T.K., Schmid–Araya, J.M., Hildrew, A.G., 2004. The contribution of subsurface
740 invertebrates to benthic density and biomass in a gravel stream. *Arch. Hydrobiol.*
741 160, 171–191.

742 Stegen, J.C., Fredrickson, J.K., Wilkins, M.J., Konopka, A.E., Nelson, W.C., Arntzen, E.V.
743 Chrisler, W.B., Chu, R.K., Danczak, R.E., Fansler, S.J., Kennedy, D.W., Resch,
744 C.T., Tfaily, M., 2016. Groundwater–surface water mixing shifts ecological
745 assembly processes and stimulates organic carbon turnover. *Nat. Commun.* 7,
746 11237.

747 Strayer, D.L., May, S.E., Nielsen, P., Wollheim, W., Hausam, S., 1997. Oxygen, organic
748 matter, and sediment granulometry as controls on hyporheic animal communities.
749 *Arch. Hydrobiol.* 140, 131–144.

750 Stubbington, R., Datry, T., 2013. The macroinvertebrate seedbank promotes community
751 persistence in temporary rivers across climate zones. *Freshwater Biol.* 58, 1202–
752 1220.

753 Stubbington, R., Wood, P.J., Boulton, A.J., 2009. Low flow controls on benthic and
754 hyporheic macroinvertebrate assemblages during supra-seasonal
755 drought. *Hydrol. Process.* 23, 2252–2263.

756 Tonina, D., Buffington, J. M. (2009). Hyporheic exchange in mountain rivers I: Mechanics
757 and environmental effects. *Geogr. Compass* 3, 1063–1086.

758 Townsend, C.R., Scarsbrook, M.R., Dolédec, S., 1997. The intermediate disturbance
759 hypothesis, refugia, and biodiversity in streams. *Limnol. Oceanogr.* 42, 938–949.

760 Valett, H.M., Hakenkamp, C.C., Boulton, A.J., 1993. Perspectives on the hyporheic zone:
761 integrating hydrology and biology. Introduction. *J. North Am. Benthological Soc.*
762 12, 40–43.

763 Vervier, P., Gibert, J., Marmonier, P., Dole-Olivier, M.J., 1992. A perspective on the
764 permeability of the surface freshwater-groundwater ecotone. *J. North Am.*
765 *Benthological Soc.* 11, 93–102.

766 Ward, A.S., 2016. The evolution and state of interdisciplinary hyporheic research. *Wiley*
767 *Interdiscip Rev* 3, 83–103.

768 Ward, A.S., Fitzgerald, M., Gooseff, M.N., Voltz, T.J., Binley, A.M., Singha, K., 2012.
769 Hydrologic and geomorphic controls on hyporheic exchange during base flow
770 recession in a headwater mountain stream. *Water Resour. Res.* 48, W04513.

771 Ward, J.V., 1989. The four-dimensional nature of lotic ecosystems. J. North Am.
772 Benthological Soc. 8, 2–8.

773 Williams, D.D., 1984. The hyporheic zone as a habitat for aquatic insects and associated
774 arthropods, in: Resh, V. H., Rosenberg, D. M. (Eds.), The Ecology of Aquatic
775 Insects. Praeger Publishers, New York, pp. 430–455.

776 Williams, D.D., Hynes, H.B.N., 1974. The occurrence of benthos deep in the substratum of a
777 stream. Freshwater Biol. 4, 233–256.

778 White, D. S., 1990. Biological relationships to convective flow patterns within stream beds.
779 Hydrobiologia 196, 149–158.

780 Wondzell, S.M., 2011. The role of the hyporheic zone across stream networks. Hydrol.
781 Process. 25, 3525–3532.

782 Wondzell, S.M., 2015. Groundwater–surface–water interactions: perspectives on the
783 development of the science over the last 20 years. Freshw. Sci. 34, 368–376.

784 Wondzell, S.M., Gooseff, M.N., McGlynn, B.L., 2010. An analysis of alternative conceptual
785 models relating hyporheic exchange flow to diel fluctuations in discharge during
786 baseflow recession. Hydrol. Process. 24, 686–694.

787 Wright–Stow, A.E., Collier, K.J., Smith, B.J., 2006. Hyporheic production is substantially
788 greater than benthic production for a common New Zealand caddisfly.
789 *Hydrobiologia* 560, 295–310.

- Yang, S., Carlson, K., 2003. Evolution of antibiotic occurrence in a river through pristine,
urban and agricultural landscapes. *Water Res.* 37, 4645–4656.
- Zeglin, H.L., 2015. Stream microbial diversity responds to environmental changes: review
and synthesis of existing research. *Front. Microbiol.* 6, 454.
- Zimmer, M.A., Lautz, L.K., 2014. Temporal and spatial response of hyporheic zone
geochemistry to a storm event. *Hydrol. Process.* 28, 2324–2337.
- Zuur, A.F., Ieno, E.N., Walker, N.J., Savaliev, A.A., Smith, G.M., 2009. Mixed Effects
Models and Extensions in Ecology with R. Springer, New York.

Legend Figures and Tables

Table 1. Glossary of terms.

Fig 1. Scheme of the streambed community structure throughout the depth profile in relation to pore size and redox potential. Arrows represent colonization depth of large macroinvertebrates (a), temporary and permanent meiofauna (b) and protozoa (c). Also shown is the theoretical boundary between benthic zone (BZ) and hyporheic zone (HZ) as the colonization limit between benthos and hyporheos. Organisms are not drawn to scale.

Fig 2. Depth-related biomass and production of invertebrates based on reported values from Smock et al. (1992) and Reynolds Jr. and Benke (2012). Predictions (black line) represent the \log_{10} biomass and production values and are derived from the linear mixed models explained in the Appendix, with 95% credible intervals (shaded grey). Open circles represent the \log_{10} -transformed values per taxa.

Fig 3. The ability of the HZ to process dissolved solutes is mediated by a hierarchical interaction between hydrological patterns and community ecology. **(a)** Daily and seasonal fluctuations between the high river-stage (H-RS) and Low river-stage (L-RS) cause variation in the open channel discharge. **(b)** The increase in the surface flow promotes higher inward water fluxes (sub-surface flow) and input of dissolved solutes into the HZ, but also the residence time of water in the sediments decreases. **(c)** Life activities of macroinvertebrates and meiofauna result in bioturbation and biorrigation phenomena in the streambed sediments, causing the occurrence of preferential flow paths and increasing permeability locally. **(d)** Protists grazing on biofilms increase its absorption surface. As a

829 result, dissolved solutes diffusion gradient is higher in presence of grazers (**d1**) than in their
830 absence (**d2**).

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

TABLE 1

| Term | Definition | Source |
|---------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|
| Hyporheic zone (Original definition) | The interstitial habitat beneath a stream, bordered by the surface water above and by the true groundwater below | Orghidan (1959) |
| Hyporheic zone (Geochemical definition) | The volume of sediment containing a specified percentage of surface water | Tonina and Buffington (2009) |
| Hyporheic zone (Hydrological definition) | The volume of sediment where water interchange between open channel and groundwater occur as a result of streambed pressure gradients and hydraulic conductivity | Tonina and Buffington (2009) |
| Hyporheic zone (Biological definition) | The volume of sediments housing a characteristic hyporheic community. This community can be defined as occasional users or permanent users | Tonina and Buffington (2009) |
| Hyporheic zone (Integrative definition) | Any location meeting four key criteria: [1] Saturate surface. [2] Existence of flow path that originate from and return to surface water. [3] Interaction with the stream occurs within a temporal scale relevant to the | Ward (2016). |

| | | |
|-------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------|
| | processes of interest. [4] Processes of interest occur continuously from subsurface to the groundwater continuum. | |
| Hyporheos | The biota occupying saturated interstitial spaces below the stream surface (benthic zone) | Standley & Boulton (1993) |
| Up–welling (UW) zone | Low–pressure areas in riverbed, where surface water comes out from HZ to the open channel | Franken et al. (2001) |
| Downwelling (DW) zone | High–pressure areas in riverbed, where surface water enter into the HZ | Franken et al. (2001) |
| Hyporheic hydrodynamics | Strength and direction of the water mass through the sediment pore spaces in the HZ, resulting from the alternation of UW and DW zones | This article |
| Micropollutants | A vast and expanding array of emerging contaminants (including pharmaceuticals, personal care products, steroid hormones, industrial chemicals and pesticides) commonly present in waters at trace concentrations, ranging from a few | Luo et al. (2014) |

| | | |
|-----------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|
| | ng/L to several µg/L. | |
| Hyporheic biorreactor | Active biological system in which the transformation of chemical compounds occurs as result of the hyporheos life activities or the active substances they produce. | This article |
| Residence time | Hydrodynamic retention time during the hyporheic exchange during which biogeochemical processesing of dissolved solutes occur in the HZ. | This article |

847

848

849

850

851

852

853

FIG 1




FIG 2




FIG 3

